

? S PN=EP 122036
S3 1 PN=EP 122036
? T S3/7

3/7/1

DIALOG(R)File 351:Derwent WPI
(c) 2005 Thomson Derwent. All rts. reserv.

004112650

WPI Acc No: 1984-258191/ 198442

Powder compsn. for nasal admin. - comprises active polypeptide and water-absorbing water- insol. base

Patent Assignee: TEIJIN LTD (TEIJ)

Inventor: NAGAI T; NAMBU N; NISHIMOTO Y; SEKINE K; SUZUKI Y

Number of Countries: 009 Number of Patents: 007

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Week
EP 122036	A	19841017	EP 84301546	A	19840308	198442 B
JP 59163313	A	19840914	JP 8337244	A	19830309	198443
US 4613500	A	19860923	US 84587779	A	19840309	198641
JP 87037016	B	19870810				198735
EP 122036	B	19890208				198906
DE 3476633	G	19890316				198912
EP 122036	B2	19941102	EP 84301546	A	19840308	199442

Priority Applications (No Type Date): JP 8337244 A 19830309

Cited Patents: 1.Jnl.Ref; AT 212; AT 329743; AT 330962; DE 1178975; DE 2229981; DE 2535258; DE 2726837; FR 2530; GB 1018125; GB 1183506; GB 1236707; GB 1461188; GB 1527605; GB 2016015; US 3155573; US 3632743; US 4294829; DE 2229987; GB 1078125; US 3682743; No-Citns.

Patent Details:

Patent No	Kind	Lan	Pg	Main IPC	Filing Notes
EP 122036	A	E	31		
Designated States (Regional): CH DE FR GB IT LI SE					
EP 122036	B	E			
Designated States (Regional): CH DE FR GB IT LI SE					
EP 122036	B2	E	17	A61K-037/02	
Designated States (Regional): CH DE FR GB IT LI SE					

Abstract (Basic): EP 122036 A

Powder compsn. for nasal admin. comprises a physiologically active polypeptide (I) or its deriv. and a water-insol. water-absorbing base.

(I) is pref. a polypeptide of molecular wt. 1000-300000, esp. calcitonin, insulin, luteinising hormone releasing hormone, desmopressin, vasopressin or oxytocin. It may be in the form of a vaccine, e.g. influenza or pertussis vaccine. Interferons may also be administered in the powder compsn. In the compsns. pref. at least 90% of the particles have a dia. of 10-250 microns.

The base may be cellulose, starch, protein, crosslinked vinyl polymer or gum. The cellulose may be cryst. cellulose, alpha-cellulose or crosslinked Na carboxymethylcellulose. The vinyl polymer is crosslinked PVP or crosslinked carboxyvinyl polymer. The water-insol. base may be used with a water-sol. water-absorbing base, e.g. lower alkyl cellulose ether, polyacrylate, polyethylene glycol or PVP.

USE/ADVANTAGE - (I) may be a high molecular wt. material readily degraded by proteolytic enzymes, so that oral admin. is ineffective and alternatives to the usual injection method are desirable.

0/2

Abstract (Equivalent): EP 122036 B

A powdery pharmaceutical composition for nasal administration comprising a physiologically active polypeptide or its derivative and a

water-absorbing and water-insoluble base

wherein at least 90 wt.% of the particles of said composition have an effective dia. ranging from 10 to 250 micras and wherein said particles do not disintegrate on spraying.

(14pp)

Abstract (Equivalent): US 4613500 A

Pharmaceutical prepn. for administration through the nose consists of (A) a physiologically active hormone, (enzyme) protein or vaccine and (B) a particulate water absorbing and water insoluble base, with at least 90 wt.% of the particles having a dia. 10-250 microns, pref. cellulose, starch, protein, crosslinked vinyl polymer or gum.

The base is pref. (a) a crystalline cellulose or (b) a crosslinked polyvinyl pyrrolidone or carboxy vinyl polymer. The base is pref. present to more than 10 times the wt. of the active substance. The hormone can be calcitonin, insulin, luteinising hormone releasing hormone, desmopressin, vasopressin or oxytocin. The vaccine is influenza or pertussive vaccine. The protein is interferon.

USE/ADVANTAGE - To avoid painful administration by injection, to provide a sustained effect; the active substance is effectively absorbed through the nasal mucosa.

(10pp)

Derwent Class: A96; B04; B07





International Patent Class (Main): A61K-037/02

International Patent Class (Additional): A61K-009/14; A61K-009/22; A61K-009/72; A61K-037/24; A61K-037/26; A61K-037/30; A61K-037/66; A61K-039/10; A61K-039/14; A61K-039/145; A61K-045/02






Powdery pharmaceutical composition for nasal administration.

Patent number: EP0122036
Publication date: 1984-10-17
Inventor: NAGAI TSUNEJI; NAMBU NAOKI; NISHIMOTO YUJI;
SEKINE KUNIO; SUZUKI YOSHIKI
Applicant: TEIJIN LTD (JP)
Classification:
- international: A61K37/02; A61K37/24; A61K37/26; A61K37/30;
A61K39/00; A61K45/02
- european: A61K9/00M14
Application number: EP19840301546 19840308
Priority number(s): JP19830037244 19830309

Also published as:

 US4613500 (A1)
 JP59163313 (A)
 EP0122036 (B2)
 EP0122036 (B1)

Cited documents:

 DE2535258
 US4294829
 US3155573
 DE2229981
 GB1236707
more >>

Abstract not available for EP0122036

Abstract of corresponding document: **US4613500**

What is provided by this invention is a powdery pharmaceutical composition for nasal administration comprising physiologically active polypeptide or its derivative such as calcitonin, insulin, etc. and a water-absorbing and water-insoluble base. The pharmaceutical composition allows polypeptide or its derivative to be efficiently absorbed through the nasal mucous membrane when it is nasally administered.

Data supplied from the **esp@cenet** database - Worldwide

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11

Publication number:

**0 122 036
A1**

12

EUROPEAN PATENT APPLICATION

21

Application number: 84301548.2

22

Date of filing: 08.03.84

51

Int. Cl.³: **A 61 K 37/02, A 61 K 37/24,
A 61 K 37/26, A 61 K 37/30,
A 61 K 39/00, A 61 K 45/02**

30

Priority: 09.03.83 JP 37244/83

43

Date of publication of application: 17.10.84
Bulletin 84/42

84

Designated Contracting States: CH DE FR GB IT LI SE

71

Applicant: **TEIJIN LIMITED**, 11 Minami
Honmachi 1-chome Higashi-ku, Osaka-shi Osaka-fu (JP)

72

Inventor: **Suzuki, Yoshiki**, 5-20-2, Tamadaira, Hino-shi
Tokyo, 191 (JP)
Inventor: **Sekine, Kunio**, 3-18-4, Tamadaira, Hino-shi
Tokyo, 191 (JP)
Inventor: **Nagai, Tsuneji**, 4-23-5, Higashiueno, Taite-ku
Tokyo, 110 (JP)
Inventor: **Nambu, Naoki**, 2-401, Nagao-jutaku 2-7,
Shibekuhoncho, Miyamae-ku Kawasaki-shi
Kanagawa-ken 213 (JP)
Inventor: **Nishimoto, Yuji**, 1-3-9, Hiratsuka,
Shinagawa-ku Tokyo, 142 (JP)

74

Representative: **Votier, Sidney David et al, CARPMAELS
& RANSFORD** 43, Bloomsbury Square, London
WC1A 2RA (GB)

54

Powdery pharmaceutical composition for nasal administration.

57

What is provided by this invention is a powdery pharmaceutical composition for nasal administration comprising physiologically active polypeptide or its derivative such as calcitonin, insulin, etc. and a water-absorbing and water-insoluble base. The pharmaceutical composition allows polypeptide or its derivative to be efficiently absorbed through the nasal mucous membrane when it is nasally administered.

EP 0 122 036 A1

POWDERY PHARMACEUTICAL COMPOSITION
FOR NASAL ADMINISTRATION

5

BACKGROUND OF THE INVENTION

Field of the Invention

10 This invention relates to a powdery pharmaceutical composition for nasal administration. More specifically, this invention relates to a powdery polypeptide or its derivative composition for nasal administration which comprises physiologically active polypeptide or its derivative such as calcitonin,
15 insulin, etc. and a water-absorbing and water-insoluble base and allows said polypeptide or its derivative to be effectively absorbed through the nasal mucosa when nasally administered.

20

Description of the Prior Art

25 Because of the fact that peptide hormones such as insulin, calcitonin, etc. have a high molecular weight and that they are readily decomposed by proteolytic enzymes such as pepsin, trypsin, chymotrypsin, etc., the peptide hormones are not absorbed sufficiently enough to display a pharmacological effect efficaciously and accordingly they have been administered by parenteral injection.

30

35 However, since the administration by injection causes pain, various attempts to develop alternative methods of administration have been made. For example, there is a method of intrarectal administration of a suppository prepared by use of such salicylic acid derivatives as sodium salicylate, 3-methoxy sodium salicylate, and 5-methoxysalicylic acid as an absorption aid (J. Pharm. Pharmacol., 33, 334,

(1981)). Besides this method, a method of intra-bronchial administration (Diabetes, 20, 552, (1971)) and a method of eye-dropping administration (Diabetes Society's Epitomes, 237, (1974)) have been studied.

5 There are, however, drawbacks to all of these methods in that they require much larger doses than injection and that their absorption is varied and accordingly any of them have hardly been put to practical use as yet.

10 On the other hand, there is a method of nasal administration of an acidic aqueous solution of insulin, wherein such a surface active agent as sodium glycocholate is used as an absorption aid, known as the attempt to develop a method of intra-
15 nasal administration (Diabetes, 27, 296, (1978)).

 However, this method may not be regarded as an expedient means, since the preparation is prepared in the form of a liquid often causing the drug to flow out of the nasal cavity upon its administration
20 into the nose and the use of a surface active agent in its preparation also causes inconvenience.

 As the powdery pharmaceutical composition for nasal administration, U.S.P. No. 4,294,829 discloses a pharmaceutical preparation comprising
25 lower alkyl ether of cellulose and a drug. This pharmaceutical composition is characterized in that its lower alkyl ether of cellulose absorbs moisture on the nasal mucous membrane, and takes the form of a viscous liquid to slowly flow over
30 the nasal mucosa and releases the drug slowly. Since the composition of this type takes the form of a viscous liquid in the nasal cavity, the drug with a high molecular weight tends to stay within the lower alkyl ether of cellulose, becoming reluc-
35 tant to the release from the composition. The composition of this type should, therefore, still undergo a significant improvement if it is to use

such a drug with a high molecular weight as calcitonin, insulin, and the like.

SUMMARY OF THE INVENTION

5

One of the objects of this invention is to provide a powdery pharmaceutical composition for nasal administration.

10

Another object of this invention is to provide a powdery pharmaceutical composition for nasal administration aimed at application of physiologically active polypeptide or its derivative.

15

A further object of this invention is to provide a powdery pharmaceutical composition for nasal administration which allows physiologically active polypeptide or its derivative to be absorbed efficiently through the nasal mucosa without the use of an absorption aid.

20

Still another object of this invention is to provide a powdery pharmaceutical composition for nasal administration which especially allows polypeptides such as insulin, calcitonin, etc. to be absorbed efficiently through the nasal mucosa without the use of an absorption aid.

25

It is yet another object of this invention to provide a powdery pharmaceutical composition for nasal administration which allows physiologically active polypeptide or its derivative to be absorbed efficiently through the nasal mucosa and also has a sustained release effect.

30

35

It is still another object of this invention to provide a powdery pharmaceutical composition for nasal administration having such a proper particle diameter as to make it possible to be administered efficiently in the nasal cavity when intranasally sprayed.

Other objects and advantages of this invention

will become apparent from the following description.

According to this invention, these objects and advantages are achieved by a powdery pharmaceutical composition for nasal administration which comprises physiologically active polypeptide or its derivative and a water-absorbing and water-insoluble base.

BRIEF DESCRIPTION OF THE DRAWING

Fig. 1 shows the decrease (%) of the plasma glucose levels after the nasal administration of a composition of this invention in which insulin is used as the polypeptide.

Fig. 2 shows the serum insulin levels after the nasal administration of a composition of this invention in which insulin is used.

As shown in Fig. 1 and Fig. 2, the powdery pharmaceutical composition for nasal administration is a composition which allows polypeptide such as insulin, etc. to be absorbed at a satisfactorily high efficiency.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

In this invention, the objective drug is physiologically active polypeptide or its derivative. As the polypeptide or its derivative, such polypeptide or its derivative with a molecular weight ranging from 1,000 to 300,000 are desirable in view of the fact that they are easily absorbed through the nasal mucous membrane. Especially those having a molecular weight ranging from 1,000 to 150,000 are more desirable.

Desirable physiologically active polypeptide or its derivative is exemplified in the following. For instance, such peptide hormones as insulin, angiotensin, vasopressin, desmopressin, felypressin,

protirelin, luteinizing hormone releasing hormone, corticotropin, prolactin, somatropin, thyrotropin, luteinizing hormone, calcitonin, kallikrein, parathyrin, glucagon, oxytoxin, gastrin, secretin, 5 serum gonadotrophin, growth hormone, erythropoietin, angiotensin, urogastrone and renin; such physiologically active proteins as interferon, interleukin, transferrin, histaglobulin, macrocortine and blood coagulation factor VIII; such enzyme proteins as lysozyme 10 and urokinase; such vaccines as acellular and cellular pertussis vaccine, diphtheria vaccine, tetanus vaccine, influenza vaccine; and such toxoids as diphtheria toxoid, tetanus toxoid and toxoids of lymphocytosis promoting factor, and filamentous hemagglutinin 15 (those are contained in acellular pertussis vaccine developed recently in Japan) may be mentioned. Of these polypeptide or its derivative mentioned above, peptide hormones, physiologically active protein and vaccine are desirable, and peptide hormones are 20 especially more desirable, and among these peptide hormones, calcitonin, insulin, luteinizing hormone releasing hormone, desmopressin, vasopressin and oxytocin are particularly desirable and calcitonin and insulin are more particularly desirable. Among 25 physiologically active protein, interferone is particularly desirable, and among vaccine, influenza vaccine and pertussis vaccine are particularly desirable.

For the preparation of powdery pharmaceutical compositions for nasal administration, the physiologically active polypeptide or its derivative is 30 preferred to be those in the form of a powder. The physiologically active polypeptide or its derivative should desirably be water-soluble in view of the fact they are to be absorbed through the nasal mucous membrane. Here, "water-soluble" means that the poly- 35 peptide or its derivative is soluble on the human nasal mucous membrane or in a similar environment,

or more concretely, it is soluble in an aqueous solution at pH around 7.4 at a temperature of about 36°C to 37°C.

5 In case where polypeptides which are water-insoluble or hardly soluble in water are to be used, it is accordingly advisable, for instance, to adjust their pH, dissolve in water, and freeze-dry, thus making them water-soluble and in the form of powder.

10 Those polypeptides which are not in the form of powder should desirably be freeze-dried into powder before use.

15 The aforementioned polypeptide or its derivative can be used in combination with human serum albumin, mannitol, sorbitol, aminoacetic acid, amino acid, sodium chloride, phospholipid, etc. for the purpose of stabilization, or for the dual purpose of stabilizing and bulking.

20 In the preparation of powdery pharmaceutical compositions for nasal administration according to the present invention, a base having duplicity properties of water-absorbing and water-insoluble is used. By the terms of water-absorbing and water-insoluble, it is meant that the base has double properties of
25 being water-absorbing and water-insoluble on the human nasal mucous membrane or in a similar environment, or more concretely, it is water-absorbing and water-insoluble at pH around 7.4 and at a temperature of 36°C to 37°C or thereabout.

30 Because of the use of a water-absorbing and water-insoluble base, the pharmaceutical composition of this invention are able to absorb the moisture on the nasal mucous membrane upon its administration into the nasal cavity, thus making each particle,
35 which is not in the state of a viscous fluid and does not flow away immediately but diffuses moderately, stay at the site on the nasal mucosa where

it adhered and allow the polypeptide or its derivative with a high molecular weight to come into thorough contact with the nasal mucosa, through which they are absorbed at a high efficiency. In
5 this invention, the selection of such a base as mentioned above for use as the base of a polypeptide or its derivative preparation for nasal application has made it possible to make the polypeptide or its derivative to be absorbed effectively and let them display their pharmacological
10 efficacy sufficiently without the use of an absorption aid.

The water-absorbing and water-insoluble bases which are to be used in this invention should be
15 distinguished from those lower alkyl ethers of cellulose, or more particularly such lower alkyl ether of cellulose as hydroxypropyl cellulose which dissolves into a viscous fluid on the nasal mucous membrane. As the desirable examples of water-absorbing
20 and water-insoluble base, the following ones may be mentioned.

They include, for instance, water-absorbing and water-insoluble celluloses such as crystalline cellulose, cellulose, α -cellulose, and cross-linked
25 sodium carboxymethyl cellulose; water-absorbing and water-insoluble starches such as hydroxypropyl starch, carboxymethyl starch, cross-linked starch, amylose, amylopectin and pectin; water-absorbing and water-insoluble proteins such as gelatin, casein, sodium
30 and casein; water-absorbing and water-insoluble gums such as gum arabic, tragacanth gum and glucomannan; and cross-linked vinyl polymers such as cross-linked polyvinyl pyrrolidone, cross-linked carboxyvinyl polymer or its salt, cross-linked polyvinyl alcohol
35 and polyhydroxyethylmethacrylate. Of these mentioned above, water-absorbing and water-insoluble celluloses and cross-linked vinyl polymer are desirable, and

water-absorbing and water-insoluble cellulose are more desirable and crystalline cellulose is especially desirable. Among cross-linked vinyl polymers, cross-linked polyvinylpyrrolidone and cross-linked carboxy vinyl polymer or its salt are desirable.

Since the quantity of a water-absorbing and water-insoluble base to be used varies depending upon the polypeptide or its derivative to be used, it can not be defined indiscriminately; however, it is desirable in general to use it in appreciable amount of more than 10 times the weight of the polypeptide or its derivative especially more than 15 times, furthermore more than 20 times.

In preparing a pharmaceutical composition of this invention, said water-absorbing and water-insoluble base may be used in combination with a water-absorbing and water-soluble base. The combined use of a water-absorbing and water-soluble base adds some degree of solubility to a water-absorbing and water-insoluble base to produce the following effect. When the pharmaceutical composition is intranasally administered, the particles which comprise a water-absorbing and water-insoluble base and polypeptide or its derivative are dispersed over the nasal mucous membrane, and the water-absorbing and water-soluble base is dissolved into the state of a viscous fluid, which gives some degree of viscosity and flowage to the whole base of this invention, thus allowing the polypeptide or its derivative to be absorbed slowly which is known as a sustained release effect.

The water-absorbing and water-soluble base may be used by simply mixing it with the water-absorbing and water-insoluble base or by mixing it with the polypeptide or its derivative at the time of their freeze-drying. In case where the water-absorbing and water-soluble base is freeze-dried together with the polypeptide or its derivative, the mixture assumes

the state in which particles of the polypeptide or its derivative are dispersed among particles of water-absorbing and water-soluble base and the final product of pharmaceutical composition comes to have much more sustained release effect.

As the water-absorbing and water-soluble base to be used in this invention, there are, for instance, polyacrylates such as sodium polyacrylate, potassium polyacrylate and ammonium polyacrylate; lower alkyl ethers of cellulose such as methyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose and sodium carboxymethyl cellulose; polyvinyl pyrrolidone, amylose, polyethyleneglycol and pullulan. Of these mentioned above, polyacrylate such as sodium polyacrylate; lower alkyl ether of cellulose such as methyl cellulose, hydroxypropyl cellulose, sodium carboxymethyl cellulose; polyethyleneglycol and polyvinyl pyrrolidone are especially desirable. The quantity of a water-absorbing and water-soluble base to be used should desirably be 0.1 to 60 wt%, or more desirably 1 to 50 wt% against a water-absorbing and water-insoluble base.

In this invention, at least 90 wt% particles of the powdery composition desirably have an effective diameter of 10 to 250 microns. By controlling the particle diameter of the particles of the powdery composition in the abovementioned range, it becomes possible to have the powdery composition diffused widely over the nasal mucous membrane when intranasally administered, make the powdery composition stay long at the place where it has adhered, and facilitate the efficient intranasal administration upon pernasal spraying of the powdery composition through the nostril.

When the powdery composition, which contains more than 10 wt% particles having an effective diameter of less than 10 microns, is nasally administered by spraying, more particles will go further

into the lungs or escape from the nostrils. Also, such powdery composition has difficulty in maintaining its effective drug concentration at a high level. On the other hand, when the powdery composition, which
5 contains more than 10 wt% particles having an effective diameter of more than 250 microns, is intranasally administered, part of the particles on the nasal mucous membrane will readily separate from there, thus undesirably making its efficacy less stayable. It is especially
10 desirable to control the powdery composition to have more than 90 wt% its particles measuring 20 to 150 microns in effective diameter.

The powdery composition of this invention may have any of the structures such as one in which the
15 water-absorbing and water-insoluble base and polypeptide or its derivative form independent particles, one in which polypeptide or its derivative particles are adhered to the surface of the water-absorbing and water-insoluble base particle, one in which polypeptide or its derivative particles are dispersed in
20 the water-absorbing and water-insoluble base particle, both forming separate phases of their own, or one in which polypeptide or its derivative particles are closely dispersed in the water-absorbing and water-insoluble base particle, thus forming a uniform
25 dispersion.

The powdery composition of this invention, which has a structure in which the water-absorbing and water-insoluble base and polypeptide or its derivative form independent particle or polypeptide or its
30 derivative particles are adhered to the surface of the water-absorbing and water-insoluble base particle, can be prepared by first adding polypeptide or its derivative to a water-absorbing and water-insoluble
35 base, then mixing them by a mechanical process, and finally filtering them to obtain the desired composition with more than 90 wt% its particles preferably

measuring 10 to 250 microns in effective diameter.

In case where a water-absorbing and water-soluble base is used in combination with a water-absorbing and water-insoluble base, both bases may be mixed together simultaneously with polypeptide or its derivatives in the mechanical mixing process.

The powdery composition of this invention, which has a structure in which polypeptide or its derivative particles are dispersed in the water-absorbing and water-insoluble base particle, both forming separate phases of their own, or polypeptide or its derivative particles are closely dispersed in the water-absorbing and water-insoluble base particle, thus forming a uniform dispersion, can be obtained by following the procedure mentioned below. It is prepared by firstly mixing polypeptide or its derivative with a water-absorbing and water-insoluble base by mechanical means, then compacting the obtained mixture under pressure, further pulverizing the compacted mixture, and finally filtering the resulting pulverulence to obtain the desired composition with more than 90 wt% its particles preferably measuring 10 to 250 microns in effective diameter. The powdery composition can also be obtained by thoroughly mixing polypeptide or its derivative with a water-absorbing and water-insoluble base in water to make a thin smooth pasty mixture, drying and pulverizing the mixture according to the ordinary methods, and finally passing the pulverized composition through a sieve.

In case where a water-absorbing and water-soluble base is to be used jointly, it may be admixed with polypeptide or its derivative and a water-absorbing and water-insoluble base in the mechanical mixing process, followed by the above-mentioned processes of compacting, etc., or a water-absorbing and water-soluble base may be

introduced into the process wherein polypeptide or its derivative is mixed with a water-absorbing and water-insoluble base in the presence of water.

There is also another method of obtaining a powdery composition by joint use of a water-absorbing and water-soluble base, wherein a water-absorbing and water-soluble base is added to polypeptide or its derivative in the process in which polypeptide or its derivative is to be freeze-dried, thus both components being freeze-dried simultaneously as mentioned above.

In order to improve the properties, appearance or odor of the powdery pharmaceutical composition of this invention, it may, if desired, contain any of known additives such as coloring agents, preservatives, antiseptics corrigents, etc. As the coloring agents, there are, for instance, β -carotene, Red No.2, Blue No.1, etc.; as the preservatives, there are stearic acid, ascorbyl stearate, ascorbic acid, etc.; as the antiseptics, there are p-hydroxybenzoate, phenol, chlorobutanol, etc., and as the corrigents, there are menthol, citrus perfume, etc.

The powdery pharmaceutical composition of this invention can be directly used as a powder for a unit dosage form.

As the desirable form for administration, the powder can be filled in capsules such as hard gelatine capsules.

As the method of pernasally applying the powdery preparation to the nasal cavity by spraying, there is, for instance, a method in which a capsule filled with the powdery preparation is placed in a sprayer, which is exclusively designed for this purpose and equipped with a needle, then the capsule is pierced with the needle to have minute holes on both the top and bottom, and thereafter jets of powder are sent into the nasal cavity by means of ballooning, etc.

The following examples illustrate the present invention more specifically; however, it should be understood that these examples are given to explain the invention and not to limit the scope of the invention.

Example 1

(i) Powdery pharmaceutical compositions for pernasal administration were obtained as follows:

(a) 400 mg of water-soluble insulin powder, which was obtained by freeze-drying a solution prepared by dissolving insulin in 0.1N-HCl aqueous solution and further adding pure water thereto, and 3,600 mg of crystalline cellulose were placed in a mixer and mixed thoroughly to obtain a uniform powdery composition, at least 90 wt% particles of which had a particle diameter of 75 to 149 microns. Thus obtained powdery composition had the insulin activity of 2.55 units/mg.

(b) 10 mg of insulin (25.5 units/mg) was dissolved in 0.2 ml of 0.1 N hydrochloric acid. 200 ml of water was added thereto to give an aqueous solution of insulin and 40 mg of polyacrylic acid (Carbopol 934) was dissolved therein. About 30 ml of an aqueous solution of 0.01 N sodium hydroxide was added to the solution to adjust it to pH 7.4. The solution was then freeze-dried to give a neutral and uniform powdery composition (I) with 5.1 units/mg of insulin, comprising insulin and sodium polyacrylate (neutral Carbopol 934).

Then, 50 mg of thus obtained powdery composition (I) and 50 mg of crystalline cellulose were put in a mortar and mixed thoroughly to obtain a uniform powdery composition (II), at least 90 wt% particles of

which had a particle diameter of 75 to 149 microns. The obtained powdery composition (II) had the insulin activity of 2.55 units/mg.

5 (d) The powdery compositions containing insulin prepared in the foregoing (a) and (b) were respectively filled in capsules to obtain insulin preparations for human pernasal administration.

(ii) The following comparative compositions were obtained to compare with the compositions of this invention.

10 (d) 700 mg of water-soluble insulin powder (25.5 units/mg), which was obtained by once dissolving insulin, then followed by freeze-drying, and 6,300 mg of lactose were placed in
15 a mixer and mixed well to give a uniform powdery composition, at least 90 wt% particles of which had a particle diameter of 75 to 149 microns. Thus obtained powdery composition contained 2.55 units/mg of insulin.

20 (e) 400 mg of water-soluble insulin powder (25.5 units/mg), which was obtained by once dissolving insulin and then freeze-drying the solution, and 3,600 mg of hydroxypropyl cellulose were placed in a mixer and mixed thoroughly
25 to give a uniform powdery composition, at least 90 wt% particles of which had a particle diameter of 75 to 149 micron. The powdery composition thus obtained contained 2.55 units/mg of insulin.

30 Example 2

(Experiment of nasal administration of powdery insulin preparation in dogs.)

3 units/kg of the respective powdery compositions of insulin prepared in Example 1, (a), (b), (d), and
35 (e) were administered intranasally to the male Beagle dogs (weighing 9.4 to 12.6 kg) which were anesthetized by intravenous injection of nembutal (containing 50 mg/ml

of pentobarbital sodium) in 25 mg/kg doses. The administration of the powdery composition of insulin was carried out by spraying it into the nasal cavity with a double balloon through a polyethylene tube (about 2 mm in diameter) inserted about 3 cm into the nostril and the blood was withdrawn from the foreleg vein as time passed after the administration. The glucose level of plasma was measured by o-toluidine (Clinical Chemistry, 8, 215 (1962)). Fig. 1 shows the decrease (%) of the plasma glucose levels from that before administration of insulin. The values shown in Fig. 1 are the average values of the four Beagle dogs. The change in plasma glucose levels after the nasal administration of 5 units/10 μ l/kg, by use of a micropipette, of an aqueous suspension of original insulin powder is shown by a broken line in Fig. 1 for the sake of comparison. In Fig. 1, (1) indicates the case where crystalline cellulose was used as the base (Example 1, (a)), (2) the case where freeze-dried insulin-sodium polyacrylate and crystalline cellulose were used (Example 1, (b)), (3) the case where lactose was used as the base (Example 1, (d)), and (4) the case where hydroxypropyl cellulose was used as the base (Example 1, (e)) respectively.

It is clear from Fig. 1 that the compositions in which crystalline cellulose is used show a high efficient absorption of insulin and that the composition in which sodium polyacrylate is used in combination with crystalline cellulose show a high efficient absorption of insulin as well as a sustained release effect.

Example 3

(i) 100 mg of porcine insulin was dissolved in 1 ml of 0.1 N hydrochloric acid, to which 40 ml of dist. water was added to obtain an insulin solution. The obtained solution was freeze-dried

to give water-soluble insulin powder (26.3 units/mg). The following powdery composition (a) of this invention was obtained from this insulin powder.

5 (a) 20 mg of the water-soluble insulin powder obtained in the above (26.3 units/mg) and 140 mg of crystalline cellulose were weighed into a mortar and mixed thoroughly to obtain a uniform powdery composition. The powdery composition thus obtained contained about 3.3 units/mg of insulin.

10 (ii) With the purpose of comparing with the preceding powdery composition (a) of this invention, the following comparative powdery compositions (b) and (c) were prepared from the aforementioned insulin powder.

15 (b) 15 mg of the water-soluble insulin powder (26.3 units/mg) and 105 mg of lactose were put in a mortar and mixed well to obtain a uniform powdery composition. The obtained powdery composition contained 3.3 units/mg of insulin.

20 (c) 20 mg of the water-soluble insulin powder (26.3 units/mg) and 140 mg of hydroxypropyl cellulose were placed in a mortar and mixed thoroughly to obtain a uniform powdery composition. The powdery composition thus obtained contained about 3.3 units/mg of insulin.

Example 4

30 (Experiment of nasal administration of powdery insulin preparation in rabbit.)

10 units/head of the respective powdery composition of insulin prepared in Example 3, (a) to (c), were administered intranasally to which native male rabbits (weighing 3.0 to 3.5 kg). The blood was withdrawn from the ear vein 10 minutes, 20 minutes, 30 minutes, and 60 minutes after the administration

and also before the administration. The administration of the powdery composition of insulin was carried out by use of a sprayer specially modified for animal use while the rabbits were lightly anesthetized by ether or not. The insulin level of the serum was determined according to the radioimmunoassay. The result is shown in Fig. 2 in which the values are the average values of the three rabbits. Also, the result, which was obtained from the experiment conducted likewise with the nasal administration of 10 units/50 μ l/head, by use of a micropipette, of an aqueous suspension of original insulin powder, is shown by a broken line in Fig. 2 for the sake of comparison.

In Fig. 2, (1) indicates the case where crystalline cellulose was used as the base (Example 3, (a)), (2) the case where lactose was used as the base (Example 3, (b)), and (3) the case where hydroxypropyl cellulose was used as the base (Example 3, (c)) respectively.

It is clear from Table 2 that the composition of this invention shows a high efficient absorption of insulin.

Example 5

2,000 mg of crystalline cellulose and 0.5 mg of freeze-dried [ASU^{1.7}]-eel calcitonin (4,000 MRC units/mg) were placed in a mortar and mixed most thoroughly to obtain a uniform powdery composition. The obtained powdery composition contained about 1 MRC unit/mg of [ASU^{1.7}]-eel calcitonin. The powdery composition was filled in the prescribed capsules with the capsule filler, each capsule containing 10 to 50 mg, to obtain a preparation for human nasal application.

Also, examples of preparation of calcitonin powdery compositions for animal experiment use are

given in the following (a) and (b).

(a) 200 mg of crystalline cellulose and 0.3 mg of freeze-dried [ASU^{1.7}]-eel calcitonin (4,000 MRC units/mg) were put in a mortar and mixed thoroughly to obtain a uniform powdery composition. The obtained powdery composition contained 5.99 MRC units/mg of [ASU^{1.7}]-eel calcitonin.

(b) 100 mg of crystalline cellulose and 0.3 mg of freeze-dried salmon calcitonin (2,000 MRC units/mg) were placed in a mortar and mixed well to obtain a uniform powdery composition. The obtained powdery composition contained 5.98 MRC units/mg of salmon calcitonin.

Example 6

(Experiment of nasal administration of powder calcitonin preparations in rabbits.)

6 MRC units/kg of the respective powdery compositions of calcitonin prepared in Example 5, (a) and (b), were administered intranasally to the white native male rabbits (weighing 2.5 to 3.0 kg). The blood was withdrawn from their ear veins before the administration and 1 hour, 2 hours, 4 hours, and 6 hours after the administration. The administration of the powdery composition was carried out in the same way as the second animal experiment made in Example 4. The concentration of the serum calcium before and after the administration was measured to study the absorbability of calcitonin through the nasal mucous membrane. The measurement of serum calcium levels was made with the use of the calcium measurement kit (IATRON Co.). The result is shown in Table 1 in which the decrease of serum calcium levels as compared with the serum calcium levels obtained before the administration of powdery preparation of calcitonin is shown in percentages. The

levels described in the table are the average values of the three rabbits.

As control, 60 MRC units/50 μ /kg of almost neutral aqueous solution of [ASU^{1.7}]-eel calcitonin was nasally administered and the result is shown in Table 1.

Table 1

	Powdery preparation		Decrease of serum calcium as compared with preadministration (%)			
	Base	Calcitonin dosage	1 hour	2 hrs	3 hrs	4 hrs
Preparation of this invention	Crystalline cellulose	[ASU ^{1.7}]-eel calcitonin 6 MRC units/kg	10.3	9.1	2.2	0.3
	Crystalline cellulose	Salmon calcitonin 6 MRC units/kg	9.6	7.5	0.7	-0.3
Control	Water	[ASU ^{1.7}]-eel calcitonin 6 MRC units/kg	2.7	1.7	0.0	0.7

Example 7

490 mg of crystalline cellulose and 10 mg of freeze-dried vasopressin (70 to 100 units/mg) were put in a mortar and mixed thoroughly to give a uniform powdery composition. The obtained powdery composition contained 1.4 to 2.0 units/mg of vasopressin.

The powdery composition thus obtained was then encapsulated according to the prescription to give a preparation for human nasal application.

Example 8

990 mg of crystalline cellulose was weighed into a mortar and 10 mg of freeze-dried luteinizing hormone releasing hormone was added thereto. They were mixed
5 thoroughly to obtain a uniform powdery composition. The obtained powdery composition contained 10 µg/mg of luteinizing hormone releasing hormone and the powdery composition was then filled in the prescribed capsules to give a preparation for human nasal appli-
10 cation.

Example 9

950 mg of crystalline cellulose was placed in a mortar, to which 50 mg of interferon (10^5 units/mg),
15 which had been freeze-dried together with human serum albumin, was added. They were mixed thoroughly to obtain a uniform powdery composition. The powdery composition thus obtained contained 5,000 units/mg of interferon and was encapsulated according to the
20 prescription to obtain a preparation for human nasal application.

Example 10

2,000 mg of crystalline cellulose and 1 mg of
25 freeze-dried desmopressin acetate were mixed thoroughly in a mortar to obtain a uniform powdery composition. The obtained powdery composition contained 0.5 µg/mg of desmopressin acetate. This composition was encapsu-
lated according to the prescription to give a prepara-
30 tion for human nasal application.

Example 11

2,000 mg of crystalline cellulose and 0.5 mg
(4,000 MRC units/mg) of freeze-dried [ASU^{1.7}]-eel
35 calcitonin or 1.0 mg (2,000 MRC units/mg) of salmon calcitonin were placed in a mortar and mixed well to obtain a uniform powdery composition. The obtained

powdery composition contained about 1 MRC unit/mg of [ASU^{1.7}]-eel calcitonin or salmon calcitonin.

10 to 50 mg of the powdery composition these obtained was then encapsulated according to the
5 prescription to give a preparation for human nasal application.

Example 12

199 mg of hydroxypropylcellulose and 1 mg (2,000
10 MRC units/mg) were dissolved in 50 ml of water. The solution was freeze-dried to give a uniform powdery composition with the salmon activity of 10 MRC units/mg, comprising salmon and hydroxypropylcellulose.

100 mg of the powdery composition and 900 mg of
15 crystalline cellulose were placed in a mortar and mixed thoroughly to obtain a uniform powdery composition, at least 90 wt% particles of which had a particle diameter of 10 to 250 microns.

Thus obtained powdery composition contained
20 1 MRC unit/mg of the salmon calcitonin. 10 to 50 mg of the powdery composition these obtained was then encapsulated according to the prescription to give a preparation for human nasal application.

25 Example 13

499 mg of hydroxypropylcellulose and 1 mg (4,000 MRC untis/mg) of [ASU^{1.7}]-eel calcitonin.

200 mg of the powdery composition and 800 mg of
30 crystalline cellulose were placed in a mortar and mixed throughly to obtain a powdery composition, at least 90 wt% particles of which had a particle diameter of 10 to 250 microns.

Example 14

35 One mg of formalin-detoxificated Partussis toxin (PT and 1 mg of formalin-treated filamentous hemagglutinin (F-HA), those were components of a

cellular pertussis vaccine developed recently in Japan and 1,000 mg of crystalline cellulose were placed in a mortar and mixed well to obtain a uniform powdery composition.

5 The obtained powdery composition contained about totally 2 µg/mg of both components.

10 to 25 mg of this composition was then encapsulated according to the prescription to give a preparation for human nasal application.

10 Thus obtained powdery composition contained 1.6 MRC units/mg of [ASU^{1.7}]-eel calcitonin. 10 to 50 mg of the powdery composition these obtained was then encapsulated according to the prescription to give a preparation for human nasal application.

15

Example 15

800 mg of hydroxypropylcellulose and 200 mg of powdery freeze-dried influenza HA vaccine were placed in a mortar and mixed well to obtain a uniform powdery composition. The powdery composition contained about 200 µg/mg of freeze-dried influenza HA vaccine. 50 mg of the powdery composition and 95 mg of crystalline cellulose were placed in a mortar and mixed well to give a uniform powdery composition, at least 90 wt% particles of which had a particle diameter of 10 to 150 microns. Thus obtained powdery composition contained 10 µg/mg of freeze-dried influenza HA vaccine. 10 to 30 mg of powdery composition these obtained was then encapsulated according to the prescription to give a preparation for human nasal application.

WHAT WE CLAIM IS:

1. A powdery pharmaceutical composition for nasal
administration comprising physiologically active
5 polypeptide or its derivative and a water-absorbing
and water-insoluble base.
2. A powdery pharmaceutical composition for nasal
administration according to Claim 1, wherein base
is any of water-absorbing and water-insoluble sub-
10 stances including celluloses, starches, proteins,
cross-linked vinyl polymers, and gums.
3. A powdery pharmaceutical composition for nasal
administration according to Claim 1, wherein base
is water-absorbing and water-insoluble celluloses.
- 15 4. A powdery pharmaceutical composition for nasal
administration according to Claim 3, wherein water-
absorbing and water-insoluble celluloses are crystal-
line cellulose, cellulose, α -cellulose, or cross-linked
sodium carboxymethyl cellulose.
- 20 5. A powdery pharmaceutical composition for nasal
administration according to Claim 3, wherein water-
absorbing and water-insoluble cellulose is crystalline
cellulose.
- 25 6. A powdery pharmaceutical composition for nasal
administration according to claim 1, wherein said
base is water-absorbing and water-insoluble cross-
linked vinyl polymers.
7. A powdery pharmaceutical composition for nasal
administration according to claim 6, wherein said
30 water-absorbing and water-insoluble cross-linked
vinyl polymers is any of cross-linked polyvinyl-
pyrrolidone and cross-linked carboxy vinyl polymer.
8. A powdery pharmaceutical composition for nasal
administration according to Claim 1, wherein water-
35 absorbing and water-insoluble base is jointly used
with a water-absorbing and water-soluble base.
9. A powdery pharmaceutical composition for nasal

administration according to Claim 8, wherein water-absorbing and water-soluble base is lower alkyl ether of cellulose, polyacrylate, polyethyleneglycol or polyvinyl pyrrolidone.

5 10. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein physiologically active polypeptide or its derivative is polypeptide or its derivative with a molecular weight of 1,000 to 300,000.

10 11. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein physiologically active polypeptide or its derivative is water-soluble polypeptide or its derivative.

15 12. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein physiologically active polypeptide or its derivative is peptide hormone, physiologically active protein, enzyme protein, or vaccine.

20 13. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein physiologically active polypeptide or its derivative is peptide hormone.

25 14. A powdery pharmaceutical composition for nasal administration according to Claim 13, wherein peptide hormone is calcitonin, insulin, luteinizing hormone releasing hormone, desmopressin, vasopressin or oxytocin.

30 15. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein physiologically active polypeptide or its derivative is vaccine.

16. A powdery pharmaceutical composition for nasal administration according to Claim 15, wherein vaccine is influenza vaccine or pertussis vaccine.

35 17. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein physiologically active polypeptide or its derivative is physiologically active protein.

18. A powdery pharmaceutical composition for nasal administration according to Claim 17, wherein physiologically active protein is interferone.

- 5 19. A powdery pharmaceutical composition for nasal administration according to Claim 1, wherein at least 90 wt% particles have an effective diameter ranging from 10 to 250 microns.

1/2

0122036

FIG. 1





FIG. 2





European Patent
Office

EUROPEAN SEARCH REPORT

0122036

EP 84301546.2

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
X	DE - A1 - 2 535 258 (FISONS LTD) * Pages 3-5, especially page 4, lines 3, 10-15; page 5, lines 23-30 *	1-3,8-10,12,17,19	A 61 K 37/02 A 61 K 37/24 A 61 K 37/26 A 61 K 37/30
	--		
D,A	US - A - 4 294 829 (YOSHIKI SUZUKI et al.) * Column 4, lines 31-64, especially lines 58-61; column 1, lines 24-38; column 1, line 53 - column 2, line 36 *	1,2,8-10,12,17,19	A 61 K 39/00 A 61 K 39/10 A 61 K 39/145 A 61 K 45/02 A 61 K 9/14 A 61 K 9/72
	--		
A	US - A - 3 155 573 (F. FOWLER) * Claims 1-3; column 5, line 47 - column 6, line 5 *	1,2,8-12,17	
	--		
A	DE - A - 2 229 981 (FISONS LTD) * Claims 1,8,9; page 3, line 30 - page 5, line 27 *	1,2,6-9,12,17,19	A 61 K 37/00 A 61 K 39/00 A 61 K 45/00
	--		
A	GB - A - 1 236 707 (STE. DE RECHERCHES INDUSTRIELLES) * Claims 1,11; page 1, lines 18-28 *	1,2,8,12,17	A 61 K 9/00
	--		
A	DE - B - 1 178 975 (RIKER LABORATORIES) * Claims 1,2,4; examples 4-6, 34; columns 6,7 *	1,2,8-14,17	
	--		
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 19-07-1984	Examiner MAZZUCCO
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	<u>US - A - 3 632 743</u> (L. GELLER et al.) * Claim 1; examples 1-6 * --	1,10-13,17	
A	<u>GB - A - 1 018 125</u> (MERCK & CO) * Example 9 * --	1,10-14,17	
A	<u>FR - M - 2 530</u> (L.-G. CHEVANCE et al.) * Claims 1,8,9; page 2, left column, paragraph 3 * --	1,10,12,15,16	
A	<u>AT - B - 330 962</u> (HOECHST) * Claims 1,2; examples 1-12 * --	1,10-14,19	TECHNICAL FIELDS SEARCHED (Int. Cl.)
A	<u>AT - B - 329 743</u> (HOECHST) * Claim; page 2, lines 14-27; examples 1,2,4-6,8 * --	1-4,8-14	
A	<u>DE - A1 - 2 726 837</u> (RECHERCHE ET INDUSTRIE THERAPEUTIQUES) * Page 5, line 5 - page 6, line 4 * --	1,10,12,15,16	
A	<u>GB - A - 1 183 506</u> (RHONE-POULENCE) * Example 7; claims 1,9,14 * --	1,10,12,15,16	
A	<u>GB - A - 1 461 188</u> (RECHERCHE ET INDUSTRIE THERAPEUTIQUES) * Page 2, line 124 - page 3, line 8 * --	1,10,12,15,16	



European Patent
Office

EUROPEAN SEARCH REPORT

0122036

EP 84301546.2

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	<p><u>AT - E - 212</u> (IOWA STATE UNIVERSITY RESEARCH FOUNDATION)</p> <p>* Page 6, lines 39-47; page 7, lines 19-39 *</p> <p>--</p>	1,2,8,10,12	
A	<p><u>GB - A - 1 527 605</u> (TAKEDA YAKUHHN KOGYO KABUSHIKI KAISHA)</p> <p>* Claim 1 *</p> <p>--</p>	1,10-14,17	
A	<p><u>GB - A - 2 016 015</u> (KEN HAYASHI-BARA)</p> <p>* Page 7, lines 28-33 *</p> <p>--</p>	1,10,12,17,18	TECHNICAL FIELDS SEARCHED (Int. Cl.)
D,A	<p>DIABETES, vol. 27, no. 3, March 1978</p> <p>SHINICHRO HIRAI et al. "Nasal Absorption of Insulin in Dogs" pages 296-299</p> <p>* Page 296, left column *</p> <p>----</p>	1,10-14,17	